

British Mathematical Olympiad 2004/5

Round 1: question 5

ANDREW JOBBINGS

www.arbelos.co.uk

12 December 2004

Question 5

Let S be a set of rational numbers with the following properties:

i) $\frac{1}{2} \in S$;

ii) if $x \in S$, then both $\frac{1}{x+1} \in S$ and $\frac{x}{x+1} \in S$.

Prove that S contains all rational numbers in the interval $0 < x < 1$.

Continued fractions

Use the notation $[a_1, a_2, \dots, a_n]$ for the continued fraction

$$\frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\dots + \frac{1}{a_n}}}}$$

where the a_i are positive integers. Note that $[a_1, a_2, \dots, a_{n-1}, 1] = [a_1, a_2, \dots, a_{n-1} + 1]$, so we may assume that $a_n \geq 2$.

To find the continued fraction of a given fraction, repeatedly invert and divide. For example, to find the continued fraction of $\frac{28}{39}$ proceed as follows:

$$\frac{28}{39} \rightarrow \frac{39}{28} = 1 + \frac{11}{28} \rightarrow \frac{28}{11} = 2 + \frac{6}{11} \rightarrow \frac{11}{6} = 1 + \frac{5}{6} \rightarrow \frac{6}{5} = 1 + \frac{1}{5} \rightarrow \frac{5}{1} = 5.$$

Hence we have $\frac{28}{39} = [1, 2, 1, 1, 5]$.

Solution to question 5

Let $f(x) = \frac{1}{x+1}$ and $g(x) = \frac{x}{x+1}$. Then $gf(0) = \frac{1}{2}$ and

$$g^{r-1}f(x) = \frac{1}{r+x}.$$

Hence

$$\begin{aligned} [a_1, a_2, \dots, a_n] &= g^{a_1-1}f \circ g^{a_2-1}f \circ \dots \circ g^{a_n-1}f(0) \\ &= g^{a_1-1}f \circ g^{a_2-1}f \circ \dots \circ g^{a_n-2}\left(\frac{1}{2}\right) \end{aligned}$$

which gives a closed form for $[a_1, a_2, \dots, a_n]$ in terms of f and g .

It follows that any rational number in the interval $0 < x < 1$ belongs to S , since any such rational may be expressed as a continued fraction of the form $[a_1, a_2, \dots, a_n]$.

Example

For example,

$$\begin{aligned} \frac{28}{39} &= [1, 2, 1, 1, 5] \\ &= f \circ g f \circ f \circ f \circ g^3 \left(\frac{1}{2} \right) \end{aligned}$$

giving the sequence of fractions

$$\frac{1}{2} \xrightarrow{g} \frac{1}{3} \xrightarrow{g} \frac{1}{4} \xrightarrow{g} \frac{1}{5} \xrightarrow{f} \frac{5}{6} \xrightarrow{f} \frac{6}{11} \xrightarrow{f} \frac{11}{17} \xrightarrow{g} \frac{11}{28} \xrightarrow{f} \frac{28}{39}.$$