


A generalisation of Pythagoras' theorem

ANDREW JOBBINGS

www.arbelos.co.uk

23 December 2008


This result was discussed in *SYMMetryplus*, Issue No 10, Autumn 1999, The proof given there made use of Hero's formula for the area of a triangle. The proof given below only makes use of the standard version of Pythagoras' theorem.



Theorem *Given a tetrahedron in which the three angles at one vertex are all right angles, as shown above, let A_1 , A_2 and A_3 be the areas of the three right-angled faces and let A be the area of the fourth triangular face. Then*

$$A^2 = A_1^2 + A_2^2 + A_3^2. \quad \square$$


Label the tetrahedron $OPQR$, where O is the vertex which has three right angles, as shown below, and let F be the foot of the perpendicular from P to QR .



Since the edge OP is perpendicular to the edges OQ and OR , it is perpendicular to the face OQR , and hence is perpendicular to OF . In other words, the triangle OFT is right-angled.

Also, OP is perpendicular to QR . Hence QR is perpendicular to both PF and OP , so it is perpendicular to OF . Therefore OF is a height of triangle OQR .

Let the lengths of four sides of the tetrahedron be a , b , c and α , as shown above, and let the perpendiculars have lengths h and H .



Now

$$\begin{aligned} A^2 &= \left(\frac{1}{2}\alpha H\right)^2 \\ &= \frac{1}{4}\alpha^2 H^2. \end{aligned}$$

Apply Pythagoras' theorem to the triangle with sides a , h and H to obtain

$$\begin{aligned} A^2 &= \frac{1}{4}\alpha^2 (h^2 + a^2) \\ &= \frac{1}{4}\alpha^2 h^2 + \frac{1}{4}\alpha^2 a^2. \end{aligned}$$

Now apply Pythagoras' theorem to the triangle with sides b , c and α to obtain

$$\begin{aligned} A^2 &= \frac{1}{4}\alpha^2 h^2 + \frac{1}{4}(b^2 + c^2)a^2 \\ &= \frac{1}{4}\alpha^2 h^2 + \frac{1}{4}a^2 b^2 + \frac{1}{4}a^2 c^2 \\ &= \left(\frac{1}{2}\alpha h\right)^2 + \left(\frac{1}{2}ab\right)^2 + \left(\frac{1}{2}ac\right)^2 \\ &= A_1^2 + A_2^2 + A_3^2. \end{aligned}$$

■